Contents

G. RICCI: Sameness between based universal algebras	3 - 22
D. DOAN, B. KIVUNGE, J. J. POOLE, J. D. H SMITH, T. SYKES, M. TEPLIT-	
SKIY: Partial semigroups and primality indicators in the fractal generation of	00.00
binomial coefficients to a prime square modulus	23 - 39
W. SIEG: Maximal classes for the family of quasi-continuous functions with closed	41 45
graph	41-45
E. STRONSKA: On \mathcal{E} -continuity and \mathcal{E} -minimality of multifunctions of two vari-	17 50
	47-52
J. SHEN, J. DONG: Existence of positive solutions to byps of higher order delay	F9 C4
C DOMAGUOWCKI Level be deside interest et al siste de la literation de la	53-04
5. DOMACHOWSKI: Local bounds and existence of solutions to non-convex dif-	65 74
M CALEWSKI. Existence of bounded solutions for fourth order Dirichlet prob	00-74
In GALE WSKI. Existence of bounded solutions for fourth order Differnet prob-	75 81
P NATH D K SINCH: On a sum form functional equation related to entropies	10-01
and some moments of a discrete random variable	83-96
O T POP M D FĂBCAS: About the bivariate operators of Durrmever-type	97-107
A. ABAL, V. GUPTA: On <i>q</i> -Baskakov type operators.	109-122
M. ŻOłDAK: On the Mazur–Ulam theorem in metric groups	123 - 130
P. KUMAM, S. PLUBTIENG: Some random fixed point theorems for random	120 100
asymptotically regular operators	131-141
H. K. PATHAK, R RODRÍGUEZ-LÓPEZ: Some results on common fixed points	
for weakly compatible mappings satisfying Altman integral type contraction.	143 - 162
P. DAS, L. K. DEY: Porosity of certain classes of operators in generalized metric	
spaces	163 - 174
S. TEMIR, G. AYKANAT: Convergence of implicit iterative process for finite	
family of <i>I</i> -nonexpansive mappings	175 - 184
N. SÖNMEZ: Division point in the Poincaré upper half plane	185 - 192
T. NOIRI, A. AL-OMARI, M. S. MD. NOORANI: Weak forms of open and closed	
functions via b - θ -open sets	193 - 203
M. PRZEMSKI: Cluster sets and related properties of multifunctions	205 - 219
J. K. KOHLI, D. SINGH: δ -perfectly continuous functions	221 - 231
W. M. MIKULSKI: Riemannian vector bundles have no canonical linear connec-	
tions	233 - 236

Gabriele Ricci

SAMENESS BETWEEN BASED UNIVERSAL ALGEBRAS

Abstract. This is the continuation of the paper "Transformations between Menger systems". To define when two universal algebras with bases "are the same", here we propose a universal notion of transformation that comes from a triple characterization concerning three representation facets: the determinations of the *Menger system, analytic monoid* and *endomorphism representation* corresponding to a basis. Hence, this notion consists of three equivalent definitions. It characterizes another technical variant and also the universal version of the very semi-linear transformations that were coordinate-free. Universal transformations allow us to check the *actual* invariance of general algebraic constructions, contrary to the seeming invariance of representation-free thinking. They propose a new interpretation of free algebras as superpositions of "analytic spaces" and deny that our algebras differ from vector spaces at fundamental stages. Contrary to present beliefs, even the foundation of abstract Linear Algebra turns out to be incomplete.

D. Doan, B. Kivunge, J. J. Poole, J. D. H Smith, T. Sykes, M. Teplitskiy PARTIAL SEMIGROUPS AND PRIMALITY INDICATORS IN THE FRACTAL GENERATION OF BINOMIAL COEFFICIENTS TO A PRIME SQUARE MODULUS

Abstract. This paper, resulting from two summer programs of Research Experience for Undergraduates, examines the congruence classes of binomial coefficients to a prime square modulus as given by a fractal generation process for lattice path counts. The process depends on the isomorphism of partial semigroup structures associated with each iteration. We also consider integrality properties of certain critical coefficients that arise in the generation process. Generalizing the application of these coefficients to arbitrary arguments, instead of just to the prime arguments appearing in their original function, it transpires that integrality of the coefficients is indicative of the primality of the argument.

Waldemar Sieg

MAXIMAL CLASSES FOR THE FAMILY OF QUASI-CONTINUOUS FUNCTIONS WITH CLOSED GRAPH

Abstract. In this paper we consider classes of functions $f : \mathbb{R} \to \mathbb{R}$. The maximal *additive* class for the family \mathcal{QU} of quasi-continuous functions with closed graph is equal to the class of all continuous functions. We also show that the maximal *multiplicative* class for \mathcal{QU} is equal to a class of continuous functions, which fulfil an extra condition.

Ewa Strońska ON E-CONTINUITY AND E-MINIMALITY OF MULTIFUNCTIONS OF TWO VARIABLES

Abstract. In this article we formulate and prove some sufficient conditions for the $l - \mathcal{E}_{X \times Y}$ continuity and the $\mathcal{E}_{X \times Y}$ - minimality of multifunctions of two variables.

Jianhua Shen, Jing Dong EXISTENCE OF POSITIVE SOLUTIONS TO BVPS OF HIGHER ORDER DELAY DIFFERENTIAL EQUATIONS **Abstract.** The paper is concerned with the existence of positive solutions for the nonlinear eigenvalue problem with singularity and the superlinear semipositone problem of higher order delay differential equations. The main results are obtained by using Guo-Krasnoselskii's fixed point theorem in cones. These results extend some of the existing literature.

Stanisław Domachowski LOCAL BOUNDS AND EXISTENCE OF SOLUTIONS TO NON-CONVEX DIFFERENTIAL INCLUSIONS

Abstract. Using a global bifurcation theorem for convex-valued completely continuous mapping we prove an existence theorem for differential inclusions of the form $u'' \in F(t, u, u')$, where F admits a convex-valued, weakly completely continuous selector and u satisfies some nonlinear boundary conditions.

Marek Galewski EXISTENCE OF BOUNDED SOLUTIONS FOR FOURTH ORDER DIRICHLET PROBLEMS WITH CONVEX-CONCAVE NONLINEARITY

Abstract. We consider the Dirichlet boundary value problem for higher order O.D.E. with nonlinearity being the sum of a derivative of a convex and of a concave function in case when no growth condition is imposed on the concave part.

P. Nath, D. K. Singh ON A SUM FORM FUNCTIONAL EQUATION RELATED TO ENTROPIES AND SOME MOMENTS OF A DISCRETE RANDOM VARIABLE

Abstract. The general solutions of a sum form functional equation have been obtained. The importance of its solutions in relation to the entropies and some moments of a discrete random variable has been discussed.

Ovidiu T. Pop, Mircea D. Fărcaş ABOUT THE BIVARIATE OPERATORS OF DURRMEYER-TYPE

Abstract. The aim of this paper is to study the convergence and approximation properties of the bivariate operators and GBS operators of Durrmeyer-type.

Ali Aral, Vijay Gupta ON q-BASKAKOV TYPE OPERATORS

Abstract. In the present paper we introduce two q-analogous of the well known Baskakov operators. For the first operator we obtain convergence property on bounded interval. Then we give the montonity on the sequence of q-Baskakov operators for n when the function f is convex. For second operator, we obtain direct approximation property on unbounded interval and estimate the rate of convergence. One can say that, depending on the selection of q, these operators are more flexible then the classical Baskakov operators while retaining their approximation properties.

Marek Żołdak ON THE MAZUR–ULAM THEOREM IN METRIC GROUPS

Abstract. Let X, Y be abelian uniquely 2-divisible groups with metrics d_X, d_Y respectively, invariant with respect to the translations and let there exist a constant c > 1 such that $d_Y(2y, 0) \ge cd_Y(y, 0)$ for $y \in Y$. We prove that each surjective isometry $U: X \to Y$ has a form U(x) = a(x) + U(0) for $x \in X$, where $a: X \to Y$ is a homomorphism.

Poom Kumam, Somyot Plubtieng SOME RANDOM FIXED POINT THEOREMS FOR RANDOM ASYMPTOTICALLY REGULAR OPERATORS

Abstract. Let (Ω, Σ) be a measurable space, X a Banach space, C a weakly compact convex subset of X and $T: \Omega \times C \to C$ a random operator. Let WCS(X) be the weakly convergent sequence coefficient of X and $\kappa_{\omega}(X)$ its Lifschitz characteristic. If T is asymptotically regular and assume that there exists $\omega \in \Omega$ and constant c such that

$$\sigma(T(\omega, \cdot)) \le c < \frac{1 + \sqrt{1 + 4WCS(X)(\kappa_{\omega}(X) - 1)}}{2}$$

we prove that T has a random fixed point. Our results also give stochastic version generalization of some results of Domínguez [Fixed point theorems for uniformly Lipschitziane mappings and asymptotically regular mappings, Nonlinear Anal. 32 No. 1 (1998), 15–27].

H. K. Pathak, Rosana Rodríguez-López SOME RESULTS ON COMMON FIXED POINTS FOR WEAKLY COMPATIBLE MAPPINGS SATISFYING ALTMAN INTEGRAL TYPE CONTRACTION

Abstract. In this paper, a common fixed point theorem for two pairs of weakly compatible mappings satisfying Altman integral type contraction in a metric space is proved. Our result extends and improves several known results.

Pratulananda Das, Lakshmi Kanta Dey POROSITY OF CERTAIN CLASSES OF OPERATORS IN GENERALIZED METRIC SPACES

Abstract. We study the porosity behavior of non-contractive mappings in a generalized metric space, a concept recently introduced in [1]. We also investigate partially the porosity position of a certain class of operators whose condition arises from [8].

Seyit Temir, Goknur Aykanat CONVERGENCE OF IMPLICIT ITERATIVE PROCESS FOR A FINITE FAMILY OF *I*-NONEXPANSIVE MAPPINGS

Abstract. We prove that an implicit iterative process converges weakly and strongly to a common fixed point of a finite family of *I*-nonexpansive mappings in a Banach space. The results presented in this paper extend and improve the corresponding results of [1, 3, 11, 12]

Nilgün Sönmez

DIVISION POINT IN THE POINCARÉ UPPER HALF PLANE

Abstract. In this work, it is shown that the coordinates of the division point can be determined by the formula in the Poincaré upper half plane.

Takashi Noiri, Ahmad Al-Omari, Mohd. Salmi Md. Noorani

WEAK FORMS OF OPEN AND CLOSED FUNCTIONS VIA b-θ-OPEN SETS

Abstract. In this paper, we introduce and study two new classes of functions called weakly b- θ -open functions and weakly b- θ -closed functions by using the notions of b- θ -open sets and b- θ -closure operator. The connections between these functions and other related functions are investigated.

Marian Przemski CLUSTER SETS AND RELATED PROPERTIES OF MULTIFUNCTIONS

Abstract. In this paper we present some types of cluster sets of multifunction. Using these concepts we relate properties of cluster sets to some generalized continuity properties, minimality of multifunctions and closedness of its graphs.

J. K. Kohli, D. Singh δ -PERFECTLY CONTINUOUS FUNCTIONS

Abstract. A new class of functions called ' δ -perfectly continuous functions' is introduced and their basic properties are studied. Their place in the hierarchy of other variants of continuity that already exist in the literature is elaborated. Further, it is shown that if X is sum connected (e.g. connected or locally connected) and Y is Hausdorff, then the function space $P_{\Delta}(X, Y)$ of all δ -perfectly continuous functions from X into Y is closed in Y^X in the topology of pointwise convergence.

W. M. Mikulski RIEMANNIAN VECTOR BUNDLES HAVE NO CANONICAL LINEAR CONNECTIONS

Abstract. We prove that Riemannian vector bundles have no canonical linear connections.